

ON THE (NON)RIGIDITY OF THE FROBENIUS ENDOMORPHISM OVER GORENSTEIN RINGS

HAILONG DAO, JINJIA LI, AND CLAUDIA MILLER

ABSTRACT. It is well-known that for a large class of local rings of positive characteristic, including complete intersection rings, the Frobenius endomorphism can be used as a test for finite projective dimension. In this paper, we exploit this property to study the structure of such rings. One of our results states that the Picard group of the punctured spectrum of such a ring R cannot have p -torsion. When R is a local complete intersection, this recovers (with a purely local algebra proof) an analogous statement for complete intersections in projective spaces first given in SGA and also a special case of a conjecture by Gabber. Our method also leads to many simply constructed examples where rigidity for the Frobenius endomorphism does not hold, even when the rings are Gorenstein with isolated singularity. This is in stark contrast to the situation for complete intersection rings. Also, a related length criterion for modules of finite length and finite projective dimension is discussed towards the end.

1. INTRODUCTION

The Frobenius endomorphism for rings of positive characteristic has been one of the central objects of study in homological commutative algebra over the past decades. Not only is it a useful tool in proofs of homological conjectures, but also its intrinsic homological properties have been shown to have strong connections with the structure of the ring or of modules over it. In this article we provide several surprising connections, for example the relationship between the ability of Frobenius to detect finite projective dimension of modules and the torsion part of the divisor class group.

First let us review some history and notations. In 1969, Kunz characterized regular local rings as those for which the Frobenius endomorphism $f: R \rightarrow R$ (or equivalently some iteration of it) is flat [23, Thm. 2.1]. Since then, a list of papers has yielded further similar homological results for f , each analogous to a classical homological result concerning the residue field k (viewed as an R -module via $\pi: R \rightarrow k$); see the survey article [28] for further details, also [5, 20]. We will use the notation ${}^f R$ for R viewed as an R -module via the n th iteration f^n of f .

For their celebrated proof of the Intersection Theorem, Peskine and Szpiro generalized one direction of Kunz's result, and shortly thereafter Herzog proved the converse, yielding the following equivalence [30, Cor. 2], [31, Thm. 1.7], [17, Thm. 3.1]:

Date: May 11, 2010.

1991 Mathematics Subject Classification. Primary: 13A35; Secondary: 13D07, 13H10.

Key words and phrases. rigidity, Frobenius endomorphism, Tor, Picard group, isolated singularity.

The third author gratefully acknowledges partial financial support from NSA grant # H98230-06-1-0035. The first author is partially supported by NSF grant 0834050.

M has finite projective dimension $\iff \mathrm{Tor}_i^R(M, {}^fR) = 0$ for all $i > 0$ and all $n > 0$

This leads one to ask to what extent the module fR could function as a test module for finite projective dimension: Is the vanishing of $\mathrm{Tor}_i^R(M, {}^fR)$ for just *one* value each of $i > 0$ and $n > 0$ sufficient? In particular, this would imply that the R -module fR is *rigid*, that is, that

$$\mathrm{Tor}_i^R(M, {}^fR) = 0 \implies \mathrm{Tor}_{i+1}^R(M, {}^fR) = 0$$

Several steps toward these goals have been made in recent years. In the general setting, Koh and Lee proved a *finiteness* result: there is a constant $c(R)$, depending only on the ring R , such that vanishing of $\mathrm{Tor}_i^R(M, {}^fR)$ for any depth $R + 1$ consecutive values of $i > 0$ and any one value of $n \geq c(R)$ implies that M has finite projective dimension [22, Prop. 2.6]. In fact, they showed that depth R consecutive values of i suffice if R is Cohen-Macaulay of positive dimension. The best possible result ([4, 13]), however, occurs in the setting of complete intersection rings:

Theorem 1.1. (*Avramov-Miller, Dutta*) *Let R be a local complete intersection and M a finitely generated R -module. Then the vanishing of $\mathrm{Tor}_i^R(M, {}^fR)$ for one value each of $i > 0$ and $n > 0$ implies that M has finite projective dimension.*

Phenomena like this can occur over non-complete intersection rings as well. In such a case, we call the corresponding fR strongly rigid (which is equivalent to being rigid when $n \geq c(R)$ by Koh and Lee's result above). See Definition 2.1 and Example 2.2) for known examples.

In Section 2, we study the properties of Gorenstein local rings whose corresponding fR is strongly rigid. We show that if R is Gorenstein such that fR is locally strongly rigid (i.e., strongly rigid at the localization at every prime ideal), then the minimal infinite projective dimension locus of a module M (see Definition 2.3) must be contained in the set of associated primes of $F^n(M)$ (see Theorem 2.5). One consequence of this result is the following characterization for modules of finite projective dimension:

Corollary 2.6 *Let R be Gorenstein local ring such that fR is locally strongly rigid for some $n > 0$ and M an R -module. Then M has finite projective dimension if and only if $\mathrm{Ass} F^n(M)$ is contained in the finite projective dimension locus of M .*

Note that the class of rings such that fR is locally strongly rigid for all $n > 0$ includes, but is strictly bigger than, the class of all local complete intersections, see Example 2.2.

We also apply Theorem 2.5 to prove that the divisor class groups of certain Gorenstein domains have no p -torsion.

Theorem 2.9 *Let R be a Gorenstein local ring such that fR is locally strongly rigid for some $n > 0$. Let I be an reflexive ideal such that I is locally free in codimension 2. Furthermore, assume that $\mathrm{Hom}_R(I, I) \cong R$. Let $q = p^n$. Then if $I^{(q)}$ satisfies Serre's condition (S_3) , I must be principal. In particular, the Picard group of the punctured spectrum of R has no p -torsion. In particular, if R satisfies condition (R_2) , $\mathrm{Cl}(R)$ has no p -torsion.*

In particular, the theorem above shows that the Picard groups of the punctured spectrum of complete intersection rings cannot have p -torsion. For complete intersections in projective spaces, such a result was first proved by Deligne in SGA 7 (Theorem 1.8, [10]) using sophisticated geometric machinery. We also note that this particular case confirms the positive characteristic case of the following conjecture we learned from [16]:

Conjecture 1.2. *(Gabber) Let $(R, \mathfrak{m})$ be a local complete intersection ring of dimension 3. Let $U_R = \operatorname{Spec}(R) - \{\mathfrak{m}\}$ be the punctured spectrum of R . Then $\operatorname{Pic}(U_R)$ is torsion-free.*

It was implied in [16] that the positive characteristic case is known, but we cannot find a precise reference. In any case, it is worth noting that our proof is purely homological and quite simple.

In Section 3, we push the ideas in the previous section further to construct many examples of Gorenstein local rings R such that $f^n R$ is not strongly rigid. In other words, the vanishing of $\operatorname{Tor}_i^R(M, f^n R)$ for just one value each of $i > 0$ and $n > 0$ is *not* sufficient to conclude that M has finite projective dimension. Two different approaches are used in these constructions. The first approach boils down to finding an isolated Gorenstein singularity with torsion class group and applying Theorem 2.9 above; see Example 3.2. The second approach takes a different route via Lemma 3.3. Obtaining an actual example requires some explicit computations on the determinantal ring of 2 by 2 minors in 9 variables and hence is less general than the first approach; see Example 3.5. The bonus is, however, that these have a torsion-free class group.

In Section 4, we study the connection between (strong) rigidity and numerical rigidity (see Definitions 4.1 and 4.2) of the Frobenius endomorphism. The main result we prove there is

Theorem 4.6 *Let R be a Cohen-Macaulay local ring with isolated singularity and of positive dimension. Fix some $n > 0$. If for every nonzerodivisor $y \in R$, $f^n(R/yR)$ is numerically rigid, then $f^n R$ is strongly rigid against modules of dimension up to one.*

The rest of the introduction contains a review of the notation and definitions used throughout the paper. We assume throughout that R is a commutative Noetherian local ring of prime characteristic $p > 0$ and that all R -modules M and N are finitely generated. The Frobenius endomorphism $f: R \rightarrow R$ is defined by $f(r) = r^p$ for $r \in R$; its self-compositions are given by $f^n(r) = r^{p^n}$. Restriction of scalars along each iteration f^n endows R with a new R -module structure, denoted by $f^n R$.

The Frobenius functor, introduced by Peskine and Szpiro in [31], is given by base change along the Frobenius endomorphism: $F_R(M) = M \otimes_R fR$ for any R -module M . Its compositions are given by $F_R^n(M) = M \otimes_R f^n R$, namely base change along the compositions f^n of f . We omit the subscript R if there is no ambiguity about R . Note particularly that the module structure on $F^n(M)$ is via usual multiplication in R on the right hand factor of the tensor product. The values of the derived functors $\operatorname{Tor}_i^R(M, f^n R)$ are similarly viewed as R -modules via the target of the base change map f^n .

It is easy to verify that $F^n(R) \cong R$ and that for cyclic modules $F^n(R/I) \cong R/I^{[q]}$, where $q = p^n$ and $I^{[q]}$ denotes the ideal generated by the q -th powers of the generators of I . For convenience, we frequently use q to denote the power p^n , which may vary.

In the sequel, we use $\ell(M)$ and $\text{pd } M$ to denote the *length* and *projective dimension*, respectively, of the module M . By codimension of M , we mean $\dim R - \dim M$. We use the notation $\mathbf{x}$ for a sequence of elements of R and often write simply $R/\mathbf{x}$ for $R/(\mathbf{x})$ to save space. Likewise, $\mathbf{x}^q$ denotes the ideal generated by the q th powers of the sequence $\mathbf{x}$, *not* the q th power of the ideal $\mathbf{x}$.

2. STRONG RIGIDITY OF FROBENIUS AND TORSION ELEMENTS IN DIVISOR CLASS GROUPS

In this section we investigate the consequences of the phenomenon that over certain rings the Frobenius map can be used to test for finite projective dimension (e.g., over complete intersection rings). This work enables us to prove strong results about torsion elements in the class groups of complete intersection rings and also allows us to construct counterexamples to such phenomena over non-complete intersection rings. We begin with some convenient definitions to facilitate the discussion.

Definition 2.1. An R -module N is called *strongly rigid* if for any integer i and any finitely generated R -module M , $\text{Tor}_i^R(M, N) = 0$ implies $\text{pd}_R M < \infty$. The module N is called *locally strongly rigid* if $N_{\mathfrak{p}}$ is strongly rigid for all $\mathfrak{p} \in \text{Spec } R$.

Example 2.2. If R is a local complete intersection ring, then $f^n R$ is locally strongly rigid for all n , see Theorem 1.1. For any local Cohen-Macaulay ring R of dimension at most 1, there is a number $c(R)$ such that for any $n \geq c(R)$, $f^n R$ is strongly rigid by virtue of Koh and Lee's result mentioned in the introduction. In particular, when $(R, \mathfrak{m})$ is Artinian and $\mathfrak{m}^{[p]} = 0$, then $f^n R$ is (locally) strongly rigid for all n , cf. [28, 2.2.8].

We also cite the following definition (see, for example, [11]):

Definition 2.3. Let M be an R -module. One defines the infinite projective dimension locus of M as

$$\text{IPD}(M) = \{\mathfrak{p} \in \text{Spec } R \mid \text{pd}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = \infty\}$$

Similarly, define $\text{FPD}(M)$ to be the finite projective dimension locus of M . Finally, we define the n -strong rigidity locus of R as

$$\text{SR}_n(R) = \{\mathfrak{p} \in \text{Spec } R \mid f^n R_{\mathfrak{p}} \text{ is strongly rigid}\}$$

The following standard facts, which we state without proof, will be used frequently:

Fact 2.4. Let $f: R \rightarrow S$ be a ring homomorphism and $\mathfrak{p}$ a prime ideal of S . Then for each $i \geq 0$ and R -module M there is a natural isomorphism

$$\text{Tor}_i^R(M, S)_{\mathfrak{p}} \cong \text{Tor}_i^{R_{f^{-1}(\mathfrak{p})}}(M_{f^{-1}(\mathfrak{p})}, S_{\mathfrak{p}})$$

Furthermore, if f is the Frobenius endomorphism of R , then $f^{-1}(\mathfrak{p}) = \mathfrak{p}$ and $R_{f^{-1}(\mathfrak{p})} \rightarrow S_{\mathfrak{p}}$ is the Frobenius endomorphism of $R_{\mathfrak{p}}$.

Theorem 2.5. Let R be a Gorenstein local ring and M an R -module. Then

$$\min \text{IPD}(M) \cap \text{SR}_n(R) \subseteq \text{Ass } F^n(M)$$

In particular, if $f^n R$ is locally strongly rigid, then

$$\min \text{IPD}(M) \subseteq \text{Ass } F^n(M)$$

Proof. Since R is Gorenstein, by the Cohen-Macaulay approximation due to Auslander and Buchweitz [3, 1.8], there is a short exact sequence

$$0 \rightarrow M \rightarrow Q \rightarrow N \rightarrow 0,$$

where $\text{pd } Q < \infty$ and N is maximal Cohen-Macaulay. Tensoring with the Frobenius endomorphism, we have an embedding

$$(2.1) \quad 0 \rightarrow \text{Tor}_1^R(N, f^n R) \rightarrow F^n(M).$$

Take any $\mathfrak{p} \in \min \text{IPD}(M) \cap \text{SR}_n(R)$; then $\text{pd}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = \infty$ and $f^n R_{\mathfrak{p}}$ is strongly rigid. It follows that $\text{pd}_{R_{\mathfrak{p}}} N_{\mathfrak{p}} = \infty$ and therefore that $\text{Tor}_1^{R_{\mathfrak{p}}}(N_{\mathfrak{p}}, f^n R_{\mathfrak{p}}) \neq 0$. On the other hand, since $\mathfrak{p}$ is minimal in the infinite projective dimension locus of M , $\text{pd}_{R_{\mathfrak{q}}} M_{\mathfrak{q}} < \infty$ for any prime $\mathfrak{q} \subsetneq \mathfrak{p}$, whence $\text{pd}_{R_{\mathfrak{q}}} N_{\mathfrak{q}} < \infty$ and so $\text{Tor}_1^{R_{\mathfrak{q}}}(N_{\mathfrak{q}}, f^n R_{\mathfrak{q}}) = 0$. Therefore, the length of $\text{Tor}_1^{R_{\mathfrak{p}}}(N_{\mathfrak{p}}, f^n R_{\mathfrak{p}})$ must be finite.

Localizing (2.1) at $\mathfrak{p}$, we have an exact sequence

$$0 \rightarrow \text{Tor}_1^{R_{\mathfrak{p}}}(N_{\mathfrak{p}}, f^n R_{\mathfrak{p}}) \rightarrow F^n(M)_{\mathfrak{p}}.$$

This implies that $\text{depth } F^n(M)_{\mathfrak{p}} = 0$. Hence $\mathfrak{p} \in \text{Ass } F^n(M)$. $\square$

Corollary 2.6. *Let R be a Gorenstein local ring such that $f^n R$ is locally strongly rigid for some $n > 0$ and M an R -module. The following are equivalent:*

- (1) M has finite projective dimension;
- (2) $\text{Ass } F^n(M) \subseteq \text{FPD}(M)$.

As an immediate consequence, we obtain the following special case with simpler hypotheses. Here, $\text{Sing}(R)$ denotes the singular locus of R . Note particularly that the hypothesis $(\min \text{Supp } M) \cap \text{Sing}(R) = \emptyset$ holds, for example, when $\dim M > \dim \text{Sing}(R)$.

Corollary 2.7. *Let R be a Gorenstein local ring such that $f^n R$ is locally strongly rigid for some $n > 0$ (e.g., if R is a local complete intersection) and M an R -module such that $(\min \text{Supp } M) \cap \text{Sing}(R) = \emptyset$. If $F^n(M)$ has no embedded primes, then M has finite projective dimension. In particular, if $F^n(M)$ is Cohen-Macaulay, then M is perfect.*

Proof. It suffices to note that

$$\text{Ass } F^n(M) = \min \text{Supp } F^n(M) = \min \text{Supp } M \subseteq \text{Spec } R \setminus \text{Sing}(R) \subseteq \text{FPD}(M)$$

where the first equality is by the assumption that $F^n(M)$ has no embedded primes, the second is well-known (see [31], for example) and the first containment follows from the hypothesis. $\square$

Remark 2.8. If R is reduced, we do not know if the condition “ $(\min \text{Supp } M) \cap \text{Sing}(R) = \emptyset$ ” in Corollary 2.7 can be replaced by the simpler condition “ $\dim M > 0$ ”. However, this is impossible when R is not reduced (see [28, 2.1.7], for example).

We now give an application of Theorem 2.5 to divisor class groups. In the sequel, we use $\text{Cl}(R)$ to denote the divisor class group of R . We refer to [15] for the definition and basic facts about $\text{Cl}(R)$ and the Picard groups and to [7] for Serre's conditions (R_n) and (S_n) .

Theorem 2.9. *Let R be a Gorenstein local ring such that fR is locally strongly rigid for some $n > 0$. Let I be an reflexive ideal such that I is locally free in codimension 2. Furthermore, assume that $\text{Hom}_R(I, I) \cong R$. Let $q = p^n$. Then if $I^{(q)}$ satisfies Serre's condition (S_3) , I must be principal. In particular, the Picard group of the punctured spectrum of R has no p -torsion. If, furthermore, R satisfies condition (R_2) , then $\text{Cl}(R)$ has no p -torsion.*

Proof. We may assume $\dim R \geq 3$. Assume that I is not principal, then it follows that $\text{pd } I = \infty$ (see [9, Corollary 11][8, Chapter VII, §4, no. 7, Corollary 2]). We claim that one can always write $I = (a) : (b)$ for $a, b \in R$. Here is a quick proof: choose a such that a generates I at the minimal primes of I . Pick an irredundant primary decomposition of (a) ; it can be written as $I \cap J$ (if $I = (a)$ we are done). Choosing b in J but not in any minimal prime of I , one can show that $I = (a) : (b)$. By the short exact sequence

$$0 \rightarrow R/(a : b) \xrightarrow{b} R/(a) \rightarrow R/(a, b) \rightarrow 0$$

we have $\text{IPD}(I) = \text{IPD}(R/(a, b))$. Thus, for any $\mathfrak{p} \in \min \text{IPD}(I)$,

$$\mathfrak{p} \in \min \text{IPD}(R/(a, b))$$

and so by Theorem 2.5,

$$\mathfrak{p} \in \text{Ass}(F^n(R/(a, b))) = \text{Ass}(R/(a^q, b^q))$$

Localize the short exact sequence

$$0 \rightarrow R/(a^q : b^q) \rightarrow R/(a^q) \rightarrow R/(a^q, b^q) \rightarrow 0$$

at $\mathfrak{p}$, and observe that $(a^q : b^q) = I^{(q)}$. From the fact that

$$\text{depth}(R/(a^q, b^q))_{\mathfrak{p}} = 0$$

we get that $\text{depth } I_{\mathfrak{p}}^{(q)} \leq 2$. On the other hand, since I is locally free in codimension 2, $\dim R_{\mathfrak{p}} \geq 3$. So, $I^{(q)}$ does not satisfy (S_3) , and our first assertion is proved. The last two statements follow immediately (note that if R is (R_2) then R is automatically normal). $\square$

As a corollary we can recover a notable result about torsion elements in the Picard groups of complete intersections.

Corollary 2.10. *Let R be an equicharacteristic local complete intersection ring of dimension at least 3. Then the Picard group of the punctured spectrum of R is torsion-free. If, furthermore, R satisfies condition (R_2) , then the class group of R is torsion-free.*

Let X be a complete intersection variety of dimension at least 2 in the projective space over a field. The Picard group of X modulo the hyperplane section is torsion-free.

Proof. Let p be the characteristic exponent of R (so it is 1 if the characteristic of R is 0). The fact that the Picard group or $\text{Cl}(R)$ has no element whose order is relatively prime to p was well-known, see [32]. Theorem 2.9 takes care of the p -torsion elements. The second half of the corollary follows by applying the first to the local ring at the origin of the affine cone over X . $\square$

Remark 2.11. The second half of the corollary was first proved by Deligne [10], and another proof was given by Bădescu [6, Theorem B]. As far as we know, ours is the first algebraic proof.

Example 2.12. *The conditions $\dim R \geq 3$ and (R_2) in the corollary cannot be weakened. Let $R = k[[x, y, z]]/(xy - z^2)$ where k is an algebraically closed field of characteristic other than 2. Then $\dim R = 2$ and R is regular in codimension 1, but $\text{Cl}(R) \cong \mathbb{Z}/(2)$ (see, for example, Proposition 11.4 of [15]).*

3. EXAMPLES OF NONRIGIDITY

In this section we construct plenty of examples of a Gorenstein ring R in positive characteristic such that fR is not (strongly) rigid. This is in stark contrast to the situation for complete intersection rings, where the strong rigidity of fR is known to hold. Our constructions take two completely different approaches. The first approach (see Example 3.2) provides the desired examples with torsion divisor class groups. This can be viewed as a natural consequence of Theorem 2.9. The second approach (see Example 3.5), on the contrary, provides the desired examples with torsion-free divisor class groups.

First we isolate a consequence of Theorem 2.9:

Corollary 3.1. *Let R be a local, Gorenstein domain with isolated singularity. Suppose that $\dim R \geq 3$ and $\text{Cl}(R)$ has a torsion element of order l that satisfies (S_3) . Then fR is not strongly rigid for any n such that $p^n \equiv 1$ or 0 modulo l . In particular, if $l = 2$, then fR is not strongly rigid for any n and not rigid for $n \gg 0$.*

Proof. Let I be a reflexive ideal which represents an l -torsion element in $\text{Cl}(R)$ and $q = p^n$. Then the ideal $J = I^{(q)}$ is isomorphic to I or R , both of which satisfy (S_3) , contradicting Theorem 2.9. When $l = 2$, for any n , $q = p^n$ is congruent to 0 or 1 modulo 2. The last statement follows from Example 2.2. $\square$

Example 3.2. *It is not hard to find examples of isolated Gorenstein singularities with torsion class group. Let $S = k[x_1, \dots, x_d]$ and l be an integer. Let T be the l -Veronese subring of S and R be the local ring at the homogeneous maximal ideal of T . Then one can show that $\text{Cl}(R) = \text{Cl}(T) = \mathbb{Z}/(l)$ using Theorem 1.6 of [34]. The ring R obviously has isolated singularity, as it is the local ring at the origin of the cone over a smooth projective variety. Also, R will be Gorenstein as long as $l \mid d$. Finally, let I represent the generator of $\text{Cl}(T)$. It is easy to see that the cyclic cover of T corresponding to I is S , so I , and therefore the generator of $\text{Cl}(R)$, is Cohen-Macaulay. In particular, it will be (S_3) . So all of the conditions of Corollary 3.1 can be satisfied easily.*

For the rest of this section we will take another approach to construct examples of non-rigidity in which the rings have *torsion-free* divisor class groups. The following result gives a general technique for finding such examples:

Lemma 3.3. *Let $(R, \mathfrak{m})$ be a Gorenstein ring with isolated singularity and positive dimension. The following are equivalent:*

- (1) fR is strongly rigid.
- (2) For any R -module L with infinite projective dimension, $\text{depth } F^n(L) = 0$

Proof of Lemma 3.3. Assume (1). Then (2) is a consequence of Corollary 2.6. Now assume (2). Let L be a module of infinite projective dimension. It is enough to prove that $\mathrm{Tor}_1^R(L, f^n R) \neq 0$. Consider the exact sequence

$$0 \rightarrow L_1 \rightarrow Q \rightarrow L \rightarrow 0$$

where Q is free and L_1 is the first syzygy of L . If $\mathrm{Tor}_1^R(L, f^n R) = 0$, then by tensoring with $f^n R$ one get

$$0 \rightarrow F^n(L_1) \rightarrow Q \rightarrow F^n(L) \rightarrow 0.$$

But since $\mathrm{pd}_R L = \mathrm{pd}_R L_1 = \infty$, one has $\mathrm{depth} F^n(L_1) = \mathrm{depth} F^n(L) = 0$. Since $\mathrm{depth} Q = \dim Q > 0$, this is a contradiction. $\square$

We also need the following crucial observation.

Lemma 3.4. *Let k be a field of characteristic $p > 0$. Let A denote the determinantal ring $k[X]/I_2$ where $X = (X_{ij})$ is a 3×3 matrix of indeterminates and I_2 is the ideal of $k[X]$ generated by all the 2×2 minors of X . Let x_{ij} denote the images of X_{ij} in A . Let $L = A/(x_{11}, x_{12})$. Then $\mathrm{depth} F^n(L) > 0$ for all $n > 0$ and $\mathrm{pd} L = \infty$.*

Proof. Let δ_{ij} denote the minors of X corresponding to X_{ij} and I be the ideal of $k[X]$ generated by X_{11}^n, X_{12}^n and all the δ_{ij} 's. We prove that for any field k (we do not need to assume that k has prime characteristic!) and any $n \geq 2$, x_{33} is a nonzerodivisor for $A/(x_{11}^n, x_{12}^n) \cong k[X]/I$. In the following paragraph, we refer the reader to [14, 15.2-4] for notations and terminologies regarding Gröbner basis.

We fix a *reverse lexicographic order* $>$ on the monomials such that

$$X_{11} > X_{12} > X_{13} > X_{21} > X_{22} > X_{23} > X_{31} > X_{32} > X_{33}$$

Using *Buchberger's Algorithm*, one can produce a Gröbner basis for I consisting of all the δ_{ij} 's, X_{11}^n, X_{12}^n and all the monomials of the form $X_{11}^l X_{12}^{n-l} X_{22}^s X_{32}^t$ where l runs through $1, 2, \dots, n-1$ and s, t run through all positive integers such that $s+t=l$. Therefore the *initial ideal* of I (henceforth $\mathrm{in}(I)$) does not contain any monomial divisible by X_{33} . Assume for some $g \in k[X]$, $X_{33}g \in I$. Let g_0 be the *remainder* of g (with respect to the generators of I) in a *standard expression* obtained by performing the *division algorithm*. If $g_0 \neq 0$, then $X_{33}g_0 \neq 0$ since $k[X]$ is a domain. On the other hand, since $X_{33}g_0 \in I$, at least one of the monomials of $X_{33}g_0$ is in $\mathrm{in}(I)$. Thus, at least one of the monomials of g_0 is in $\mathrm{in}(I)$. This contradicts the fact that g_0 is a nonzero *remainder*. Thus $g_0 = 0$ and $g \in I$. It follows that x_{33} is a nonzerodivisor for $k[X]/I$.

Finally, we show $\mathrm{pd} L = \infty$. Assume that $\mathrm{pd} L < \infty$, i.e., the ideal (x_{11}, x_{12}) is of finite projective dimension. By a result of MacRae [26, Cor. 4.4], two-generated ideals of finite projective dimension have the form $a(b, c)$, where a is a nonzerodivisor and b, c form a regular sequence. But if $(x_{11}, x_{12}) = a(b, c)$ for such a, b and c , since the degree of x_{11} is one, a is forced to be a unit (otherwise, (x_{11}, x_{12}) would be a principal ideal which is impossible). Therefore $(x_{11}, x_{12}) = (b, c)$. But since $x_{11}x_{22} - x_{21}x_{12} = 0$, (x_{11}, x_{12}) cannot be an ideal generated by a regular sequence of two elements. This is a contradiction. $\square$

Combining the two lemmas above, we get the following example. Note that the divisor class group of the ring in this example is isomorphic to $\mathbb{Z}$ [7, 7.3.5], which is torsion-free.

Example 3.5. Let R be the localization of the determinantal ring A as in Lemma 3.4 with respect to the maximal ideal (X) . Then fR is not strongly rigid for any n .

Remark 3.6. In view of Koh-Lee's theorem mentioned in the Introduction, Example 3.5 immediately yields the nonrigidity of fR for any $n \geq c(R)$ (cf. Example 2.2). But in fact, with a little further computation, the reader can check that this example yields nonrigidity for *all* $n > 0$: indeed, the module N of infinite projective dimension constructed in Theorem 2.5 by taking for the module M the module L of Lemma 3.4 satisfies $\mathrm{Tor}_1^R(N, {}^fR) = 0$ by the argument in the proof, but it can be shown that in fact $\mathrm{Tor}_2^R(N, {}^fR) \neq 0$.

We point out that we do not know of any example showing that fR is not (strongly) rigid when $\dim R = 0$ or against a module M of finite length. See, however, the discussion at the end of Section 4.

4. SOME FURTHER OBSERVATIONS

Throughout this section, d will always be the dimension of the ring and n always denotes some positive integer. We know from the previous section that R could fail to be strongly rigid when the ring R is no longer complete intersection. However, we still hope that to some extent such a property could hold over non-complete intersection rings. In particular, we do not know any example showing that fR is not rigid when $\dim R = 0$ or against a module M of finite length.

We first make two more definitions, the first of which is just a refinement of the definition of strongly rigidity of fR .

Definition 4.1. Let h be a nonnegative integer. fR is called *strongly rigid against modules of dimension at most h* , if for any integer i and any finitely generated module M of dimension at most h , $\mathrm{Tor}_i^R(M, {}^fR) = 0$ implies $\mathrm{pd}_R M < \infty$.

Definition 4.2. fR is called *numerically rigid* if for any R -module M of finite length, $\ell(F^n(M)) = p^{nd}\ell(M)$ implies $\mathrm{pd}_R M < \infty$.

The second definition above is motivated by the following characterization for modules of finite projective dimension and finite length over complete intersection rings.

Theorem [12, 28] *Let R be a complete intersection ring in characteristic p and M an R -module of finite length. Then the following are equivalent:*

- (1) M has finite projective dimension,
- (2) $\ell(F^n(M)) = p^{nd}\ell(M)$ for all $n > 0$,
- (3) $\ell(F^n(M)) = p^{nd}\ell(M)$ for some $n > 0$.

The implication (3) $\Rightarrow$ (1) in the above theorem simply says that if R is a complete intersection ring, then fR is *numerically rigid* for any n . When R is no longer a complete intersection ring, it is an open question whether fR could still be numerically rigid¹. In fact, such a question is closely related to the rigidity question discussed earlier in this paper. The goal of this section is to explore the connections between them.

¹the implication (1) $\Rightarrow$ (2) in the theorem fails even over Gorenstein rings, see [29]

The following technical result plays a crucial role here. Recall that if $\ell(M \otimes N) < \infty$ and $\text{pd } N < \infty$, then $\chi(M, N) \stackrel{\text{def}}{=} \sum_{j=0}^{\text{pd } N} (-1)^j \ell(\text{Tor}_j^R(M, N))$.

Proposition 4.3. *Let R be a Noetherian local Cohen-Macaulay ring of positive dimension and of characteristic $p > 0$. Let M be an R -module of codimension c . Suppose $\dim M > 0$ and $R_{\mathfrak{p}}$ is a complete intersection ring for every minimal prime $\mathfrak{p}$ of M . Then*

$$\ell(F_{R/\mathbf{x}}^n(M/\mathbf{x}M)) \geq q^c \chi(M, R/\mathbf{x}), \forall n > 0,$$

for any s.o.p. $\mathbf{x}$ of $F^n(M)$ which is also R -regular. Moreover, equality holds for some $n > 0$ if and only if: (1) $F^n(M)$ is Cohen-Macaulay and (2) for every minimal prime $\mathfrak{p}$ of M , $\text{pd}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} < \infty$.

For the proof the properties of the higher Euler characteristics of Koszul complexes are used in an essential way. We recall some terms and results here.

For a pair of modules M, N such that $\ell(M \otimes N) < \infty$ and $\text{pd } N < \infty$, the i th higher Euler characteristic is defined by the following formula

$$\chi_i(M, N) = \sum_{j=i}^{\text{pd } N} (-1)^{j-i} \ell(\text{Tor}_j^R(M, N)).$$

By convention, $\chi(M, N) = \chi_0(M, N)$. Some standard facts about χ and χ_i can be found in [25, 33]. In this paper, we particularly need the following two well-known results due to Serre and Lichtenbaum [25, Lemma 1 and Theorem 1], [33].

Lemma 4.4. *Let M be an R -module and $\mathbf{x} = \{x_1, x_2, \dots, x_c\}$ an R -sequence such that $\ell(M/\mathbf{x}M) < \infty$. Then $\chi(M, R/\mathbf{x}) \geq 0$, with the equality holding if and only if $\dim M < c$.*

Theorem 4.5. *Let M be an R -module and $\mathbf{x}$ an R -sequence such that $\ell(M/\mathbf{x}M) < \infty$. Then for any $i > 0$, $\chi_i(M, R/\mathbf{x}) \geq 0$, with the equality holding if and only if $\text{Tor}_i(M, R/\mathbf{x}) = 0$ (and hence $\text{Tor}_j(M, R/\mathbf{x}) = 0$ for all $j \geq i$).*

Proof of Proposition 4.3. Since $\text{Supp } M = \text{Supp } F^n(M)$ (see [31]), $\min \text{Supp } F^n(M) = \min \text{Supp } M$. We have

$$\begin{aligned} \ell(F_{R/\mathbf{x}}^n(M/\mathbf{x}M)) &= \ell(F^n(M) \otimes_R R/\mathbf{x}) \\ &\geq \chi(F^n(M), R/\mathbf{x}) \\ &= \sum_{\mathfrak{p} \in \min \text{Supp } M} \ell(F^n(M)_{\mathfrak{p}}) \chi(R/\mathfrak{p}, R/\mathbf{x}) \\ &= \sum_{\mathfrak{p} \in \min \text{Supp } M} \ell(F_{R_{\mathfrak{p}}}^n(M_{\mathfrak{p}})) \chi(R/\mathfrak{p}, R/\mathbf{x}) \\ &\geq \sum_{\mathfrak{p} \in \min \text{Supp } M} q^c \ell(M_{\mathfrak{p}}) \chi(R/\mathfrak{p}, R/\mathbf{x}) \\ &= q^c \chi(M, R/\mathbf{x}) \end{aligned}$$

where the first inequality holds since $\chi_1(F^n(M), R/\mathbf{x}) \geq 0$ by Theorem 4.5, the second and fourth equalities by Lemma 4.4, and the second inequality is a result over complete intersection rings [12, Thm. 1.9] (note that $R_{\mathfrak{p}}$ is complete intersection by the hypotheses).

Therefore, furthermore, equality holds if and only if $\chi_1(F^n(M), R/\mathfrak{x}) = 0$ and $\ell(F_{R_{\mathfrak{p}}}^n(M_{\mathfrak{p}})) = q^c \ell(M_{\mathfrak{p}})$ for every minimal prime $\mathfrak{p}$ of M . The former is equivalent to $F^n(M)$ being Cohen-Macaulay by Theorem 4.5 and the latter is equivalent to $M_{\mathfrak{p}}$ having finite projective dimension over $R_{\mathfrak{p}}$ by [28, Thm. 5.2.2] since $R_{\mathfrak{p}}$ is complete intersection. $\square$

Theorem 4.6. *Let R be a Cohen-Macaulay local ring with isolated singularity and of positive dimension. Fix some $n > 0$. If for every nonzerodivisor $y \in R$, ${}^fR/yR$ is numerically rigid, then fR is strongly rigid against modules of dimension at most one.*

Proof. Let M be an R -module of dimension at most one. Assume ${}^fR/yR$ is numerically rigid for every nonzerodivisor $y \in R$. We want to prove that for any $i > 0$, $\text{Tor}_i(M, {}^fR) = 0$ implies $\text{pd } M < \infty$. Let $\mathbf{x} = \{x_1, \dots, x_{d-1}\}$ be an R -sequence contained in $\text{Ann } M$. We may assume that $i = 1$ by replacing M by its $(i - 1)$ th syzygy over the ring $R/(x_1, \dots, x_{d-1})$ and using that $\text{Tor}_i(R/(x_1, \dots, x_{d-1}), {}^fR) = 0$ for all $i > 0$ as $\text{pd}_R R/(x_1, \dots, x_{d-1}) < \infty$.

Letting K be the first syzygy of M as an $R/(x_1, \dots, x_{d-1})$ -module, we get a short exact sequence

$$0 \rightarrow F^n(K) \rightarrow F^n((R/(x_1, \dots, x_{d-1}))^t) \rightarrow F^n(M) \rightarrow 0.$$

It follows that $F^n(K)$ is a Cohen-Macaulay module of dimension one. Hence by Proposition 4.3 (note that R has isolated singularity), one has $\ell(F_{R/yR}^n(K/yK)) = q^{d-1} \chi(K, R/yR)$ for every $y \in R$ which is regular on both K and R . Therefore, $\ell(F_{R/yR}^n(K/yK)) = q^{d-1} \ell(K/yK)$. Since we assume ${}^fR/yR$ is numerically rigid, K/yK has finite projective dimension over R/yR . Thus K has finite projective dimension over R , whence M does too by the long exact sequence of Tors against the residue field k . $\square$

Remark 4.7. For the determinantal ring $R = k[X]/I_2$ used in Section 3, it was shown there that fR is not strongly rigid against modules of dimension at most 5 for any n . In fact, we can also modify the example a little bit to show that it is not strongly rigid against modules of dimension at most 3. For k of arbitrary characteristic though, we do not know if fR is strongly rigid against modules of dimension at most 0, 1, or 2. However, in characteristic 2 we have an example which shows that fR is not strongly rigid against modules of dimension 1. In fact, if we set $k = \mathbb{Z}/2\mathbb{Z}$ and take the module $N = R/(x_{12}, x_{13}, x_{21}, x_{23}, x_{31}, x_{32})$, then it is easy to check that $\dim N = 1$, $\text{depth } F(N) = 1$ and $\text{pd } N = \infty$. Taking an R -sequence x_1, x_2, x_3, x_4 contained in the annihilator of N and embedding N into a module of finite projective dimension over $R/(x_1, x_2, x_3, x_4)$ (via the Auslander-Buchweitz short exact sequence again), the cokernel of this embedding gives such an example. Therefore, by Theorem 4.6, we also obtain an example of Gorenstein ring R in characteristic 2 for which the corresponding R -module fR is not numerically rigid.

ACKNOWLEDGEMENTS

We thank Craig Huneke for crucial remarks which connected our works and Srikanth Iyengar who carefully read an earlier version of this manuscript and suggested a better statement for Corollary 2.6. We also thank the anonymous referee for many helpful comments.

REFERENCES

1. I. Aberbach, J. Li, *Asymptotic vanishing conditions which force regularity in local rings of prime characteristic*, Math. Res. Lett. **15** (2008), 815–820.
2. M. Auslander, M. Bridger, *Stable module theory*, Mem. Amer. Math. Soc. **94** (1969).
3. M. Auslander, R. Buchweitz, *The homological theory of maximal Cohen-Macaulay approximations*, Colloque en l'honneur de Pierre Samuel (Orsay, 1987), Mem. Soc. Math. France (N.S) **38** (1989), 5–37.
4. L. L. Avramov, C. Miller, *Frobenius powers of complete intersections*, Math. Res. Lett. **8** (2001), 225–232.
5. L. L. Avramov, S. Iyengar, C. Miller, *Homology over local homomorphisms*, Amer. J. Math. **128** (2006), 23–90.
6. L. Bădescu, *A remark on the Grothendieck-Lefschetz theorem about the Picard group*, Nagoya Math. J. **71** (1978), 169–179.
7. W. Bruns, J. Herzog, *Cohen-Macaulay Rings*, Cambridge Studies in Adv. Math. **39**, Cambridge University Press, Cambridge, 1993.
8. N. Bourbaki, *Elements of Mathematics. Commutative Algebra*, Hermann, London, 1972.
9. A. Braun, *On a question by M. Auslander*, J. Algebra, **276**, No. 2 (2004), 674–684.
10. P. Deligne, *Cohomologie des intersections complètes*, Sem. Geom. Alg. du Bois Marie (SGA 7, II), Springer Lect. Notes Math., No. 340, (1973).
11. H. Dao, *Decency and rigidity over hypersurfaces*, arXiv math.AC/0611568, preprint
12. S. P. Dutta, *Frobenius and multiplicities*, J. Algebra **85** No. 2 (1983), 424–448.
13. S. P. Dutta, *On modules of finite projective dimension over complete intersections*, Proc. Amer. Math. Soc. **131** No. 1 (2003), 113–116.
14. D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics, **150**, Springer-Verlag, New York, 1995.
15. R. M. Fossum, *The divisor class group of a Krull domain*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band **74**, Springer-Verlag, New York-Heidelberg, 1973.
16. O. Gabber, *On purity for the Brauer group*, Arithmetic Algebraic Geometry, Oberwolfach Report No. 34 (2004), 1975–1977.
17. J. Herzog, *Ringe der Charakteristik p und Frobenius-Funktor*, Math. Z. **140** (1974), 67–78.
18. C. Huneke, R. Wiegand, *Tensor products of modules and the rigidity of Tor*, Math. Ann. **299** (1994), no. 3, 449–476.
19. C. Huneke, R. Wiegand, *Correction to: “Tensor products of modules and the rigidity of Tor”*, Math. Ann. **338** (2007), no. 2, 291–293.
20. S. Iyengar, S. Sather-Wagstaff, *G -dimension over local homomorphisms. Applications to the Frobenius endomorphism*, Illinois J. Math. **48** (2004), 241–272.
21. I. Kaplansky, *Commutative Rings*, revised edition. The University of Chicago Press, Chicago, Ill.-London, 1974.
22. J. Koh, K. Lee, *Some restrictions on the maps in minimal resolutions*, J. Algebra **202** (1998), 671–689.
23. E. Kunz, *Characterization of regular local rings for characteristic p* , Amer. J. Math. **91** (1969), 772–784.
24. J. Li, *Characterizations of regular local rings in positive characteristics*, Proc. Amer. Math. Soc. **136**, (2008), 1553–1558.
25. S. Lichtenbaum, *On the vanishing of Tor in regular local rings*, Illinois J. Math. **10** (1966), 220–226.
26. R. E. MacRae, *On the homological dimension of certain ideals*, Proc. Amer. Math. Soc. **14** (1963), 746–750.
27. C. Miller, *A Frobenius characterization of finite projective dimension over complete intersections*, Math. Z. **233** (2000), 127–136.
28. C. Miller, *The Frobenius endomorphism and homological dimensions*, Contemp. Math. **331** (2003)
29. C. Miller, A. K. Singh, *Intersection multiplicities over Gorenstein rings*, Math. Ann. **317** (2000), 155–171.
30. C. Peskine, L. Szpiro, *Sur la topologie des sous-schémas fermés d'un schéma localement noethérien, définis comme support d'un faisceau cohérent localement de dimension projective finie*, C. R. Acad. Sci. Paris Sér. A-B **269** (1969), A49–A51.

31. C. Peskine, L. Szpiro, *Dimension projective finie et cohomologie locale*, Inst. Hautes Etudes Sci. Publ. Math. **42** (1973), 323-395.
32. L. Robbiano, *Some properties of complete intersections in “good” projective varieties*, Nagoya Math. J. **61** (1976), 103-111.
33. J. P. Serre, *Algèbre Locale. Multiplicités*, Lecture Notes in Mathematics, vol. 11, Springer-Verlag, Berlin/Heidelberg/New York, 1975.
34. K.-i. Watanabe, *Some remarks concerning Demazure’s construction of normal graded rings*, Nagoya Math. J. **83** (1981), 203-211.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS, LAWRENCE, KS 66045-7523 USA

E-mail address: `hdao@math.ku.edu`

DEPARTMENT OF MATHEMATICAL SCIENCES, MIDDLE TENNESSEE STATE UNIVERSITY, MURFREESBORO, TN 37132 USA

E-mail address: `jinjiali@mtsu.edu`

MATHEMATICS DEPARTMENT, SYRACUSE UNIVERSITY, SYRACUSE, NY 13244-1150 USA

E-mail address: `clamille@syr.edu`